



# CCA ADVANTAGE

The Voice of the Certified Crop Adviser Program  
[www.agronomy.org/cca](http://www.agronomy.org/cca)

## CCAs Play Important Role in Asian Soybean Rust Indemnity



By Luther Smith  
CAE, Executive  
Director — CCA  
ph: 608/268-4977  
e-mail: [lsmith@agronomy.org](mailto:lsmith@agronomy.org)

**U**SDA's Risk Management Agency (RMA) has issued a news release recommending that producers document they followed good farming practices as advised by local agricultural experts, such as extension agents and certified crop consultants, when combating Asian soybean rust. (Editors note: certified crop consultants include certified crop advisers and certified professional agronomists.)

Under the terms of the Common Crop Insurance Policy, a practice is considered a good farming practice if agricultural experts agree that the production method used will allow the crop to make normal progress toward maturity and produce at least the yield used to determine the production guarantee. The failure to purchase and apply adequate control measures for Asian soybean rust due to economic reasons is not an insurable cause of loss.

If crops become infected, RMA recommends that insured producers document:

- the date of discovery of the disease,
- any recommendations received from agricultural experts and
- actions taken regarding the application of appropriate control measures.

As with all crop insurance policies and plans of insurance, farmers must

use good farming practices. This ensures that in the event of any naturally occurring disease outbreaks, such as soybean rust, producers will be eligible for an indemnity based on the full amount of the loss. If good farming practices are not followed, production attributed to the failure to follow good farming practices is assessed, resulting in a reduction in the indemnity.

For more information regarding good farming practices and crop insurance protection against Asian soybean rust, please see the crop policies area on the RMA Web site at <http://www.rma.usda.gov>.

Additional links to Web sites with information on Asian soybean rust can be found on the American Society of Agronomy Web site at [www.agronomy.org](http://www.agronomy.org).

### Should Something Be Done About Ethics?



By Kim Polizotto  
ICCA Standards and  
Ethics Committee Chair  
ph: 317/462-2485  
e-mail: [polizotto@potashcorp.com](mailto:polizotto@potashcorp.com)

**W**isconsin has revoked a CCA certification due to falsifying CEU records — primarily having someone sign someone else's name at a CEU event.

Northeast Region Certified Crop Adviser Chair Jeff Ten Eyck wrote in a recent column, "I must remind everyone about the seriousness of falsifying continuing education unit (CEU) sign-in sheets. It is a

breach of the Code of Ethics we all signed and it is simply not acceptable."

ICCA Board Chair Steve Dlugosz wrote a column in February CCA Advantage, "Ethics — Doing the Right Thing." He stated, "I've heard from individuals who are frustrated that a few CCAs are doing things like this and are still in the program."

Several participant comments in the recently completed CCA Program Survey stressed the importance of getting tougher with ethics.

If you are one of those CCAs who feel it is time for your local board to take a stronger position on ethics breaches, start by reviewing the CCA Code of Ethics, Article V, Duty to the Profession.

*continued on page 19*

### ICCA OFFICERS

#### CHAIRMAN

**Stephen Dlugosz**  
Agrilience  
ph: 317/432-5562  
[stevedlugosz@comcast.net](mailto:stevedlugosz@comcast.net)

#### VICE-CHAIRMAN

**Fred Vocasek**  
Servi-Tech  
ph: 620/227-7123  
[fredv@servi-techinc.com](mailto:fredv@servi-techinc.com)

#### PAST-CHAIRMAN

**Robert Beck**  
Pioneer Hi-Bred International, Inc.  
ph: 515/334-6778  
[robert.beck@pioneer.com](mailto:robert.beck@pioneer.com)

#### EXECUTIVE DIRECTOR

**Luther Smith**  
American Society of Agronomy  
ph: 608/268-4977  
[lsmith@agronomy.org](mailto:lsmith@agronomy.org)



## ICCA Board Working for You

**T**he International Certified Crop Adviser Board of Directors met in Washington, DC, March 21-23, 2005, for their spring board meeting. On Tuesday, March 22, they took advantage of being in our nation's capital to conduct briefing sessions for House and Senate Ag Committee staffers.

Explaining the purpose and role of the CCA Program were Board Chair Steve Dlugosz (Agrilience, IL), Fred Vocasek (Servi-Tech, KS), and Past Chair Bob Beck (Pioneer Hi-Bred Intl., IA). After the formal presentation, board members spoke informally with the legislative staffers to answer questions and reiterate the points made during the presentation.

Following the briefing, board members called on their own legislators to familiarize them with the CCA program and to leave contact information for future reference. This was the first opportunity many board members had to make Hill visits and they are looking forward to expanding the event in February 2006. It is expected that the next farm bill will be in the process of development at that time and there will be plenty for ICCA Board members to talk to legislators about.



ICCA Board members and guests at the spring meeting, Washington, DC.



**Russell Duncan**, South Carolina CCA (left), and **Tom Kemp**, CCA southern regional representative, discuss the Certified Crop Adviser Program with **Christy Seyfert** of Sen. Chambliss's office.



(L to r) **Steve Dlugosz**, CCA chairman; **Jane Shey**, DC consultant; **Bob Beck**, CCA past-chairman; and **Dave Sleper**, president-elect ASA, discuss policy issues.

Other Board actions include:

- Directed the chair to appoint an ad hoc committee to evaluate all the program fees and recommend a course of action.
- Adopted a set of procedures for the development of CEU specialty areas.
- Voted to become a 501c6 organization.



(L to r) **Luther Smith**, CCA executive director; **Bob Stephenson**, USDA's FSA representative to the CCA Board; and **Steve Dlugosz**, CCA chairman



## Chairman's Corner

# I'm Excited



By Steve Dlugosz, CCA  
Chairman International CCA Program  
Agronomist, Agriliance, Indianapolis, IN  
ph: 317/432-5562  
e-mail: [stevedlugosz@comcast.net](mailto:stevedlugosz@comcast.net)

**I** really am! Your International Board met earlier this spring in Washington, DC, for its semi-annual meeting. We have been working hard to increase our visibility in DC, and to get the CCA Program better connected with key policy makers in agriculture. I would like to briefly share some successes.

The first day the board met with representatives from three government agencies.

**FSA** (Farm Service Agency) asked for our input on how it can more effectively utilize the private sector in delivering technical assistance to the farm. This is a great opportunity to show FSA how CCAs work with farmers every day, and how we can assist with the proper implementation of government programs on the farm. Appropriate funding for private-sector efforts is critical, and several of our CCA Board members will be concentrating on this.

**EPA** made a presentation on water quality trading and the role of the private sector in this endeavor. Again it asked for CCAs' input and assistance. Several board members volunteered to work directly with EPA on this project.

**NRCS** also gave an update. The CCA Program has enjoyed a long relationship with this agency and will continue to work closely with it as new initiatives are developed and implemented.

The CCA Program is rapidly building a reputation for working well with different agencies in Washington and is increasingly recognized as being a trusted adviser on the farm. I really enjoyed the remark North Central Representative Dave Franzen made: "It used to be, 'Hello, I'm from the government, I'm here to help.' Now it's, 'I'm from the government, can you help us?'" This change in attitude has the potential to be a great opportunity for many CCAs across the country.

The board spent the next day on Capitol Hill telling the story of who we are and what we do. We met with staffers from the House Agriculture Committee in the morning, then with staffers from the Senate Ag Committee in the afternoon. We made three short presentations — what is a certified crop adviser, how CCAs affect farmer profitability and an example how CCAs will play a front-line role in combating Asian rust. Our message was well received and we answered a number of very good questions. Individual board members also made personal visits to their respective Senate and congressional offices.

### WHY WORRY ABOUT WASHINGTON?

You may be wondering why all the effort in DC? Well, a new farm bill is starting to be developed. We continue to hear that government agencies will not be adding additional staff and will need to utilize more of the private sector to implement future programs. CCAs are very qualified to be a large part of that private sector. Now is the time to work hard so everyone recognizes and understands the CCA Program. We must convince them that CCAs are well positioned to assist with these efforts, *with the proper compensation*. We're on the farm every day. We have the expertise. We have our farmers' trust.

Increasing recognition of the CCA Program and building relationships in Washington remind me of trying to establish a good stand of alfalfa. They take a pretty good investment of time, effort and money. I hope the fruits of our labor produce some high-quality hay for CCAs in the years to come.

## Should Something Be Done About Ethics? *(continued from page 17)*

### Duty to the Profession:

1. An applicant shall aid in exclusion from certification those who have not followed this Code of Ethics or who do not have the required education and experience.
2. An applicant shall uphold this Code of Ethics by precept and example and encourage, by counsel and advice, other registrants to do the same.

3. An applicant having positive knowledge of deviation from this code by another registrant shall bring such deviation to the attention of the registrant's local board.

An ethics inquiry starts with you. It is up to you to take a stand. It is up to you to write a letter of concern to your board. The hardest part of the process is signing your name, but without your name on the letter it can go nowhere. It may be one of the hardest things you have ever done, and it won't be taken lightly. You are a professional.



# Tillage and Poultry Litter Application Effects on Cotton Growth and Yield

By Chandra K. Reddy, E. Z. Nyakatawa, and D.W. Reeves

### EARN ONE CEU!

All CCAs may earn up to 20 Continuing Education Units (CEUs) per two-year cycle as board-approved self-study articles which will include CCA Advantage articles. The CCA CEU logo (above) marks all pre-approved material, with the CEU value indicated by the number in the middle. To receive one CEU in nutrient management, read this article, fill out the attached exam and mail the tear-out form, along with \$10, to the American Society of Agronomy.

**A**doption of no-tillage cotton production in the southern U.S. increased from about 254,000 ha in 1998 to 784,000 ha in 2002. Conservation-tillage cotton acreage nearly tripled in Alabama and Georgia during this period. A survey by the National Cotton Council of America reported that 57% of the total cotton hectareage in the southeast U.S. was under no-tillage, which resulted in an average savings of \$50.03 per ha (\$20.13 per acre) for fuel and labor compared with conventional tillage.

Problems reported with no-till cotton include soil compaction, poor seedling emergence, poor plant establishment, stunted growth and reduced yields. Cotton does not produce enough residues to supply the C necessary to increase soil organic matter and improve soil tilth in the seed zone. Also, cotton residues do not last long after harvest to protect the soil from erosion and reduce loss of soil moisture from evaporation. Therefore, without additional residues to supplement cotton residues, soils under no-till cotton may develop a crust at the surface and a compacted layer in the top 5 to 10 cm.

Inclusion of winter cover crops in no-till cotton systems can provide crop residues to make conservation-tillage cotton production systems comply with the standards set by the Natural Resources Conservation Service. Benefits include improving soil water retention, increasing soil organic matter and reducing erosion. Winter cover crops may also reduce nitrate leaching to groundwater by picking up excess nutrients remaining from the summer cotton crop.

Crop rotations of different genus or species improve soil fertility, reduce erosion, reduce the buildup of pests and increase net profits. Corn can be grown as a summer crop in rotation with cotton to break the life cycles of major cotton insect pests and diseases. Corn also supplies additional residues to increase soil organic matter in conservation-tillage cotton production systems. Cotton, corn and winter rye, which are dicot, monocot and monocot respectively, have root systems that complement

each other in nutrient uptake when grown in rotation, making them more efficient in using soil nutrients. This may reduce buildup of excess nutrients such as P, which is associated with application of poultry litter based on N content.

Application of poultry litter as a source of N and P has been shown to increase yields of crops such as corn and pastures. Furthermore, our studies have shown that poultry litter improves soil chemical properties compared with inorganic sources of N such as ammonium nitrate. Application of poultry litter to cotton will provide an environmentally sustainable way of disposing of the large quantities of waste in this region.

The objective of this study was to evaluate the effect of tillage systems on growth and yield of cotton grown in rotation with a winter rye cover crop with poultry litter and ammonium nitrate fertilizer application in north Alabama. A field study was conducted at the Alabama Agricultural Experiment Station, Belle Mina, AL, on a Decatur silt loam soil from 1996 to 2001. Treatments consisted of three tillage systems: conventional till, mulch till and no-till; two cropping systems: cotton in summer and fallow in winter and cotton in summer and rye in winter; three N levels: 0, 100 and 200 kg N ha<sup>-1</sup>; and two N sources: ammonium nitrate and poultry litter. Ammonium nitrate was used at one N rate (100 kg N ha<sup>-1</sup>), which is the recommended rate for cotton in the Tennessee Valley region.

Poultry litter was broadcast by hand and incorporated to a depth of 5 to 8 cm by preplant cultivation in conventional tillage and mulch tillage systems, whereas in the no-tillage system, it was not incorporated. The ammonium nitrate and poultry litter were applied to the plots 1 day before cotton planting. Before planting, the plots received a blanket application of 336 kg ha<sup>-1</sup> of a 0-20-20 fertilizer each year from 1996 to 1999, 112 kg ha<sup>-1</sup> of a 0-0-60 fertilizer in 2000, and 224 kg ha<sup>-1</sup> of 5-20-20 fertilizer in 2001 and 2002 to minimize the effects of P and K applied through poultry litter.

### RESULTS AND DISCUSSION

The years 1998, 2000 and 2001 had poor rainfall distribution for cotton. The years 1998 and 2000 were characterized by droughts in May or June and/or July while 2001 had excess rainfall over the same period.

**Winter Rye and Surface Residue Cover.** In 1998 winter rye biomass yield in plots that had received 100 and 200 kg N ha<sup>-1</sup> in the form of poultry litter under cotton was 112% and 130% greater than in plots that received 100 kg N ha<sup>-1</sup> in the form of ammonium nitrate, respectively. Similar figures in 2000 were 14% and 68% greater, and in 2001, 50% and 150% greater, respectively. Cumulative winter rye cover crop biomass yields



## Continuing Education Self-Study Course

### Nutrient Management

due to application of 100 kg N ha<sup>-1</sup> in the form of ammonium nitrate, 100 kg N ha<sup>-1</sup> in the form of poultry litter, and 200 kg N ha<sup>-1</sup> in the form of poultry litter were 4,329, 5,402 and 7,638 kg ha<sup>-1</sup>, respectively.

The above data show that poultry litter application to cotton has more residual positive effects on the amount of biomass produced by the winter rye cover crop compared with ammonium nitrate when used at the same rate of 100 kg N ha<sup>-1</sup>. The significance of these results is that since the winter rye crop is grown without additional fertilizer, it can scavenge residual N from the poultry litter, which would otherwise be susceptible to leaching during the winter and spring. The winter rye cover crop may also reduce sediment loss of P from the plots by tying P in plant biomass during the winter when there is no cotton.

There was a significant year x tillage x cropping system interaction on surface residue cover estimated immediately after cotton planting. Surface residue cover immediately after cotton planting in conventional tillage with winter rye cover cropping was 20% and 13% in 1997 and 1998, respectively, compared with 1% in conventional tillage with winter fallow. Similar values for 2000 and 2001 were 36% and 34%, respectively, in conventional tillage with winter rye cover cropping compared with an average of 5% in conventional tillage with winter fallow cropping.

There was a significant tillage x N source interaction on surface residue cover immediately after cotton planting. In the conventional tillage system, where crop residues were incorporated into the soil, application of 100 kg N ha<sup>-1</sup> in the form of poultry litter increased surface residue cover to 30% compared with 23% for 100 kg N ha<sup>-1</sup> in the form of ammonium nitrate. However, in mulch tillage and no-tillage systems, where crop residues were either partially incorporated into the soil or not incorporated at all, there was no improvement in surface residue cover due to application of poultry litter compared with ammonium nitrate. This was expected since a greater proportion of the soil surface would already be covered with crop residues under mulch tillage and no-tillage systems and should not be taken to imply that application of poultry litter did not increase the amount of residues left on the soil surface.

Our results suggest that surface application of poultry litter instead of ammonium nitrate can offer further protection to the soil against erosion in a conventional tillage system.

**Cotton Seedling Emergence and Establishment.** Inadequate cotton seedling emergence and establishment and consequently variable crop stands have been blamed for poor adoption of conservation tillage in cotton production for the southeastern U.S. There was a significant year x tillage system interaction on cotton seedling counts. Cotton seedling counts under conventional tillage averaged over cover cropping systems and N treatments were similar to those under no-tillage and mulch tillage systems in each year of study. The rate of emergence in no-till was significantly greater than that in conventional tillage. Also, in plots that received 100 kg N ha<sup>-1</sup> in the form of poultry litter and 200 kg N ha<sup>-1</sup> in the form of poultry litter, rate of seedling emergence was significantly greater than in plots that did not receive N and plots that received 100 kg N ha<sup>-1</sup> in the form of ammonium nitrate in all years. This was attrib-

uted to higher volumetric soil moisture content in the top 7 to 10 cm of the soil. The optimum number for cotton seedling establishment is about 10 plants m<sup>-2</sup>. Our results show that final cotton seedling counts were in this optimum range in 1997 and 2001. In 1998 and 2000, when soil moisture was most limiting during seedling emergence, surface residue cover was positively correlated with final cotton seedling counts, which in turn were positively correlated to leaf area index, number of bolls per plant, biomass and lint yield.

### COTTON GROWTH AND YIELD PARAMETERS

**Plant Height** — There was a significant year x tillage system and year x N source interaction on cotton plant height. In 1997, plant height under no-tillage was 10 cm greater than under conventional tillage. This can be attributed to the fact that no-tillage improved cotton growth by conserving soil moisture during the drought period of July 1997. Cotton plant height for plants that received 100 kg N ha<sup>-1</sup> in the form of ammonium nitrate was 20, 12 and 15 cm greater than plants in plots that received 100 kg N ha<sup>-1</sup> in the form of poultry litter in 1997, 1998 and 2001, respectively. However, in 2000 there were no differences in plant height between plants that received 100 kg N ha<sup>-1</sup> in the form of ammonium nitrate and those that received 100 kg N ha<sup>-1</sup> in the form of poultry litter, which may indicate poultry litter was able to compensate for lower nutrient availability by conserving soil moisture during the dry spells of May and July. Plant height significantly correlated with number of bolls per plant, biomass yield and lint yield over the 4-year period, indicating plant height is a good indicator of cotton productivity.

**Leaf Area Index** — There was a significant tillage x cropping system and year x N source interaction on cotton leaf area index at full bloom. In conventional tillage plots, cotton leaf area index was 5.80 with winter rye cover cropping compared with 4.80 without winter rye cover cropping. However, in no-till with winter rye cover cropping, cotton leaf area index was 5.30, which was only 0.2 units higher compared with winter fallow cropping. Cotton following winter rye had higher leaf area index compared with cotton after winter fallow, but differences were not significant. Also, cotton leaf area index for cotton winter rye cropping system under conventional tillage was 0.5 units higher than that under no-till. Leaf area index for plants that received 100 kg N ha<sup>-1</sup> in the form of ammonium nitrate was 1.90, 1.00 and 1.40 units greater than for plants that received 100 kg N ha<sup>-1</sup> in the form of poultry litter in 1997, 1998 and 2001, respectively. Leaf area index is a good indicator of plant growth and soil conditions for plant productivity, and it positively correlated with bolls per plant, biomass yield and lint yield.

**Number of Bolls per Plant** — There was significant tillage x cropping system, tillage x N source and year x tillage interaction on number of cotton bolls per plant. In no-tillage system, winter rye cover cropping increased the number of bolls per plant by 7 compared with cotton winter fallow cropping, which had 21 bolls per plant. However, in conventional tillage, winter rye cover cropping did not have a significant effect on bolls per plant. Without rye cover cropping, no-tillage had slightly lower



## Continuing Education Self-Study Course

### Nutrient Management

bolts per plant compared with conventional tillage. However, with rye cover cropping, no-till had, on average, 6 more bolts per plant compared with conventional tillage, showing that rye cover cropping was essential to the reproductive development of cotton under no-tillage system.

In mulch tillage and no-till plots, plants that received 100 kg N ha<sup>-1</sup> in the form of ammonium nitrate had 9 and 8 more bolts per plant compared with plants that received 100 kg N ha<sup>-1</sup> in the form of poultry litter, respectively. No-tillage system had 4 and 2 more bolts per plant compared with conventional tillage system in 1997 and 2001, respectively. Number of bolts per plant positively correlated with cotton biomass yield and lint yield.

**Lint Yield** – There was significant year x N source and tillage x N source interaction on cotton lint yield. In 1998 cotton lint yield in plots that received 100 kg N ha<sup>-1</sup> in the form of ammonium nitrate averaged 1,536 kg ha<sup>-1</sup>, which was 19%, 15% and 34% greater than lint yields in 1997, 2000 and 2001, respectively. In plots that received 100 kg N ha<sup>-1</sup> in the form of poultry litter, the highest lint yield was 1,354 kg ha<sup>-1</sup> in 1998, which was 17% and 15% greater than lint yields in 1997 and 2001. This variation in yield responses in each year can be explained in terms of rainfall distribution during May, June and August.

There were no significant differences in cotton lint yield between 100 kg N ha<sup>-1</sup> ammonium nitrate and 100 kg N ha<sup>-1</sup> poultry litter treatments in plots under conventional tillage system. However, for mulch tillage and no-tillage systems, plants in plots that received 100 kg N ha<sup>-1</sup> ammonium nitrate had 12% and 11% higher lint yield compared with plots that received 100 kg N ha<sup>-1</sup> poultry litter, respectively. In plots that received 100 kg N ha<sup>-1</sup> in the form of ammonium nitrate, lint yield in mulch tillage and no-tillage systems was 7% and 13% greater than in conventional tillage. However, in plots that received 100 kg N ha<sup>-1</sup> in the form of poultry litter, lint yield in conventional tillage system was 12% greater than under mulch tillage system and 4% greater than under no-till. These results can be attributed to the fact that soil incorporation of poultry litter under conventional tillage speeds up mineralization whereas in mulch tillage and no-tillage systems, poultry litter mineralization is slower. In no-tillage with winter rye cropping, lint yields averaged about 100 kg ha<sup>-1</sup> greater than in no-tillage with winter fallow cropping during the same period. Compared with conventional tillage with winter fallow cropping, lint yields in no-tillage with winter rye cropping averaged 137 kg ha<sup>-1</sup> higher during the study period.

The key to increasing cotton lint yields is using conservation tillage (mulch-till or no-till) with adequate N fertility and soil moisture in the critical stages of cotton growth and development; namely, seedling emergence, squaring, flowering and boll development to maturity. These stages include the months of May, June, July and August. Benefits of conservation tillage are mainly a result of keeping crop residues on the soil surface, which improves the plant environment by holding additional moisture. This will further improve soil organic matter and reduce soil erosion.

Breaking up and incorporation of crop residues during tillage leaves little or no residue on the surface. Therefore, the benefits of cover cropping such as reduction in surface evapo-


ration and erosion control are diminished. In addition, crop residue incorporation results in immobilization of inorganic N, which affects early plant growth. Tillage promotes oxidation of crop residues and soil organic matter, which are important in soil moisture conservation. Therefore, for benefits of cover cropping to be realized, crop residues need to be left intact on the soil surface to reduce soil moisture evaporation and also to slow down the rate of decomposition. Without winter rye cover cropping, no-tillage with 100 kg N ha<sup>-1</sup> gave similar or slightly lower yields compared with conventional tillage with the same N rate of 100 kg N ha<sup>-1</sup>.

Nitrogen application in the form of ammonium nitrate or poultry litter significantly increased cotton lint yield in conventional tillage except for the 100 kg N ha<sup>-1</sup> in the form of poultry litter treatment in 1997. In mulch tillage plots where poultry litter was incorporated into the soil, there were no significant differences in lint yields between the 100 kg N ha<sup>-1</sup> in the form of ammonium nitrate and the 100 kg N ha<sup>-1</sup> in the form of poultry litter treatments in all years. With 200 kg N ha<sup>-1</sup> of poultry litter and cotton winter rye cover cropping, lint yields under no-tillage were up to 28% (or 351 kg N ha<sup>-1</sup>) greater than under conventional tillage with 100 kg N ha<sup>-1</sup> of ammonium nitrate and winter rye cover cropping. However, with 100 kg N ha<sup>-1</sup> of poultry litter, no-till did not do better than conventional tillage with 100 kg N ha<sup>-1</sup> in the form of ammonium nitrate, which further supports the need for adequate N fertilization in no-tillage.

#### SUMMARY

Mulch tillage and no-tillage systems did not have adverse effects on cotton seedling emergence and establishment compared with conventional tillage, contrary to previous reports. Generally, cotton growth parameters in plots that received the same rate of N in the form of ammonium nitrate were better than those that received poultry litter. However, during drought years, no-tillage compensated for reduced availability of N from poultry litter by conserving soil moisture. In the no-tillage system, winter rye cover cropping significantly increased number of cotton bolts compared with winter fallow cropping. Use of no-tillage without a cover crop in cotton production may not give significant benefits. Rainfall distribution in May, June, July and August had significant effect on cotton lint yields. Similarly to what was observed with cotton growth parameters, application of poultry litter at the rate of 100 kg N ha<sup>-1</sup> generally gave lower or similar cotton lint yield compared with ammonium nitrate at the same rate, whereas at 200 kg N ha<sup>-1</sup>, lint yields were significantly greater than those at 100 kg N ha<sup>-1</sup>, irrespective of the N source.

**Editor's note:** Content was adapted from the paper "Tillage and Poultry Litter Application Effects on Cotton Growth and Yield," which was published in *Agronomy Journal*, Vol. 96, November-December 2004, and is courtesy of the authors Chandra K. Reddy, E. Z. Nyakatawa, and D.W. Reeves.



## Continuing Education Self-Study Course

### Nutrient Management

#### Get a CEU!

This exam is worth 1 CEU in **Nutrient Management**. An exam score of 70% or higher will earn CEU credit. The International CCA program has approved self-study CEUs for 20 of the 40 CEUs required in the two-year cycle.

#### DIRECTIONS

1. Read the self-study article on pages 26-28 carefully.
2. Answer the questions by clearly marking an "X" in the box next to the best answer for each question.
3. Complete the self-study exam registration form on the back of this page.
4. Clip out this self-study examination page, fold and place in envelope.
5. Enclose a check for \$10.00 made payable to the American Society of Agronomy, for processing fees. Payment in U.S. funds only.
6. **Mail your self-study exam and fee to:**  
ASA c/o CCA Self-Study Exam, 677 S. Segoe Road, Madison, WI 53711. *Please allow 60 days for processing.*
7. An electronic version of this test is also available at [www.AgProfessional.com](http://www.AgProfessional.com). Go to the Certified Crop Advisers section (lefthand column) and access the "CCA Advantage" link.

## Tillage and Poultry Litter Application Effects on Cotton Growth and Yield

### May/June Self-Study Examination

1. During the period from 1998 to 2002, no-till cotton acreage in the southern U.S.  
 a. decreased.  
 b. stayed the same.  
 c. doubled.  
 d. tripled.
2. Seedling emergence may be compromised in no-till continuous cotton because of  
 a. soil crusting.  
 b. heavy residue cover.  
 c. nitrogen tie-up.  
 d. allelopathy.
3. A reason to use a winter rye cover crop in a no-till cotton production system is to  
 a. increase soil organic matter.  
 b. increase residual soil nitrates available to following crops.  
 c. break cotton insect and disease cycles.  
 d. eliminate herbicide carryover.
4. This research study of tillage and poultry litter application included  
 a. four nitrogen levels.  
 b. two nitrogen sources.  
 c. two types of poultry litter.  
 d. three site/years.
5. Growth of a winter rye cover crop in this study was highest using a nitrogen source of  
 a. poultry litter.  
 b. ammonium nitrate.  
 c. anhydrous ammonia.  
 d. diammonium phosphate.
6. Cotton plant height is a good indicator of  
 a. lint yield.  
 b. lint quality.  
 c. tillage system used.  
 d. boll size.
7. At 100 kg N ha<sup>-1</sup> applied as poultry litter, the highest yields came from  
 a. subsurface banding.  
 b. mulch tillage.  
 c. no-till.  
 d. conventional tillage.
8. As compared to ammonium nitrate treatments in conventional tillage, lint yields using poultry litter in conventional tillage were  
 a. similar.  
 b. higher.  
 c. lower.  
 d. much lower.



DETACH HERE



# Continuing Education Self-Study Course


## Nutrient Management

9. As compared to cotton after winter fallow, the leaf area index of cotton following winter rye was

- a. significantly lower.
- b. slightly lower.
- c. slightly higher.
- d. the same.

10. The most important months for cotton to benefit from adequate soil moisture include

- a. April, May, June, and July.
- b. May, June, July, and August.
- c. July, August, and September.
- d. August, September, and October.



### SELF-STUDY EXAM REGISTRATION FORM

Name: \_\_\_\_\_

Address: \_\_\_\_\_

City: \_\_\_\_\_ State/Province: \_\_\_\_\_ Zip: \_\_\_\_\_

CCA Certification #: \_\_\_\_\_

Credit Card #: \_\_\_\_\_ Type of Card: Visa  Mastercard  Discovery  Am Express

Expiration Date \_\_\_\_\_ Name on Card: \_\_\_\_\_

Enclose a \$10 check payable to American Society of Agronomy.

X

Signature of Registrant as it appears on Code of Ethics

I certify that I alone completed this self-study course and recognize that an ethics violation may revoke my CCA status.

**This exam issued June 2005 expires June 2008.**

DETACH HERE

### SELF-STUDY EXAM EVALUATION FORM

Rating Scale: 1=Poor 5=Excellent

Information presented will be useful in my daily crop advising activities: 1 2 3 4 5

Information was organized and logical: 1 2 3 4 5

Graphics/tables were appropriate and enhanced my learning: 1 2 3 4 5

I was stimulated to think how to use and apply the information presented: 1 2 3 4 5

This article addressed the stated competency area and performance objective(s): 1 2 3 4 5

Briefly explain any "1" ratings: \_\_\_\_\_

Topics you would like to see addressed in future self-study materials: \_\_\_\_\_